

MITOGEN-ACTIVATED SIGNALING AND CELL CYCLE REGULATION IN AIRWAY SMOOTH MUSCLE

Kristen Page and Marc B. Hershenon

Department of Pediatrics, University of Chicago Children's Hospital, 5841 S. Maryland Avenue, MC 4064, Chicago, IL 60637-1470

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Brief review of cell cycle regulation
4. Growth factor stimulation of airway smooth muscle cells
5. Role of ERK signaling pathway in airway smooth muscle cell cycle progression
6. Role of PI 3-kinase
7. Role of Rac1
8. Inhibition of airway smooth muscle cell proliferation
9. Potential role of protein kinase C isoforms
10. Perspective
11. Acknowledgements
12. References

1. ABSTRACT

Increased airway smooth muscle mass has been demonstrated in patients with bronchopulmonary dysplasia and asthma. These data highlight the need for a precise understanding of the events involved in airway smooth muscle mitogenesis. To that end, investigators have developed cell culture systems adopting tracheal and bronchial myocytes from different species. A growing body of literature suggests that common signal transduction pathways regulate airway smooth muscle cell cycle entry across species lines. This review summarizes what is known about mitogen-activated signal transduction in airway smooth muscle cells. The extracellular signal regulated kinase (ERK) and phosphatidylinositol 3-kinase (PI 3-kinase) pathways appear to be major positive regulators of airway smooth muscle proliferation. It is also conceivable that growth factor stimulation of airway smooth muscle simultaneously elicits signaling through negative regulatory pathways such as the p38 mitogen-activated protein (MAP) kinase pathway, perhaps as a safeguard against excessive growth.

2. INTRODUCTION

Growing airways are covered with a well-formed layer of smooth muscle cells by the end of the embryonic period of fetal lung development (1). With airway circumferential and axial growth, this layer increases in size, a result of both cellular hypertrophy and hyperplasia. It has been demonstrated that smooth muscle mass is abnormally increased in the airways of premature infants with bronchopulmonary dysplasia (2, 3), due in part to excess cell proliferation (4). Data describing the growth of airway smooth muscle during normal postnatal development are lacking, however.

Increased airway smooth muscle mass has also been demonstrated in patients with non-fatal (5) and fatal asthma (5-13). In the most convincing study to date, Ebina and colleagues (13) examined the airway thickness and smooth muscle cell number of patients with fatal asthma using state-of-the-art stereological techniques. Two subgroups of asthmatic airways were found, one in which smooth muscle mass was increased only in the central bronchi (Type I) and another in which smooth muscle thickness was increased throughout the airway tree (Type II). In Type I, smooth muscle hyperplasia was responsible for central airway smooth muscle thickening, whereas in Type II, cellular hypertrophy was present over the length of the airway tree.

Increased expression of epidermal growth factor (EGF), a mitogen for human airway smooth muscle (14), has been noted in the airways of asthmatic patients (15, 16). In addition, when added to cultured human airway smooth muscle cells, bronchoalveolar lavage fluid from asthmatic airways increased the ERK activation, cyclin D₁ protein abundance, [³H]-thymidine incorporation and cell number, relative to that from control subjects (17). Finally, excess airway smooth muscle DNA synthesis has been demonstrated in two animal models of airways disease, hyperoxic exposure and allergen sensitization (18-20).

Taken together, the above data strongly suggest that excess smooth muscle proliferation is present in the airways of patients with bronchopulmonary dysplasia and asthma, and highlight the need for a precise understanding of the events involved in airway smooth muscle mitogenesis. To that end, investigators have developed cell culture systems adopting tracheal and bronchial myocytes

Airway myocyte cell cycle regulation

from different species. A growing body of literature suggests that common signal transduction pathways regulate airway smooth muscle cell cycle entry across species lines. A summary of mitogen-activated signal transduction in airway smooth muscle follows below.

3. BRIEF OVERVIEW OF CELL CYCLE REGULATION

Regulation of mammalian cell proliferation by extracellular signals occurs in the transition from G_0 to G_1 of the cell cycle. A key event in this transition is phosphorylation of the 110 kD retinoblastoma protein (Rb) by the cyclin D₁/cyclin-dependent kinase-4 (cdk4) dimer. Cyclin D₁, cdk4, proliferating cell nuclear antigen (PCNA) and a cyclin-dependent kinase inhibitor, p21^{Cip1}, are induced as part of the delayed early response to mitogenic stimulation (21). The cyclin D₁/cdk4 dimer titrates p27^{Kip1}, another inhibitor of cdk activity (22), and also enters into complexes with PCNA and p21^{Cip1} (23, 24). PCNA is a cofactor of DNA polymerase delta (25). p21^{Cip1} may act as an assembly factor, promoting binding of cyclin D₁ with cdk4. As G_1 progresses, p27^{Kip1} is degraded by the ubiquitin-proteasome pathway (26). Once enough cyclin D₁ and cdk4 are synthesized and enough p27^{Kip1} degraded, steric inhibition by p27^{Kip1} is exceeded, leading to phosphorylation and activation of cdk4 by cdk4-activating kinase (CAK) (27, 28). Activation of cdk4, in turn, leads to hyperphosphorylation of Rb (29), a key regulator of S-phase traversal. Rb is phosphorylated near the G₁/S restriction point and accumulates phosphate until cells exit from mitosis (30). Once phosphorylated, Rb releases the transcription factors E2F1-3, which activate genes required for DNA replication such as DNA polymerase alpha (31). Titration of p27 by cyclin D₁/cdk4 also frees cyclin E/cdk2 complexes, which contribute to Rb phosphorylation and the modification of preinitiation complexes which trigger DNA replication (32). Finally, in late G_1 , cyclin A is induced and assembles with cdk2, marking the irreversible decision to enter S phase (G₁ restriction point) (33).

Ectopic overexpression of either cyclin D or cyclin E accelerates progression through G and reduces the proliferative requirement for serum-derived growth factors (34-36). Conversely, abolition of cyclin D or cyclin E activity through the use of neutralizing antibodies or antisense oligonucleotides effectively blocks entry into S phase (34, 37, 38). The requirement of cyclin D₁ for S phase traversal has been confirmed in bovine tracheal myocytes (39).

4. GROWTH FACTOR STIMULATION OF AIRWAY SMOOTH MUSCLE CELLS

Many studies have focused on smooth muscle cell proliferation in response to growth factor stimulation. Airway smooth muscle cells proliferate in response to peptide growth factors ligating receptor tyrosine kinases (40-43) as well as to bronchoconstrictor substances associated with G protein-coupled seven transmembrane receptors (44-51). Sensitivity to growth factor activation

appears to be species-specific. For instance, histamine is mitogenic for human airway smooth muscle (51, 52), but does not induce proliferation in bovine cells (41). Nevertheless, the signaling pathways and downstream targets initiated by growth factor stimulation appear to be remarkably constant across species lines (see below).

For the last five years, our laboratory has concentrated on the signaling pathways responsible for PDGF-induced DNA synthesis in bovine tracheal myocytes. We will therefore focus on this system as a model of airway smooth muscle proliferation. The PDGF receptor comprises two distinct chains (A and B) which are dimerized by a disulfide bond, and can exist in three possible isoforms, AA, AB, or BB. Typically, growth factor binding to a receptor tyrosine kinase activates the receptor's intracellular kinase domain, leading to the phosphorylation of specific tyrosine residues inside the kinase domain. Ligand binding of PDGF induces formation of a stable receptor dimer, which in turn causes one receptor molecule to phosphorylate the other in the dimer (53). EGF, a monomeric molecule, likely binds simultaneously to two receptor molecules (54). Phosphorylation of tyrosine residues inside the kinase domain further increases kinase activity, leading to phosphorylation of other sites in the receptor outside the kinase domain (55). These phosphotyrosine residues serve as binding sites for downstream signal transduction molecules containing Src-homology 2 (SH2) domains. For example, the PDGF receptor holds nine phosphotyrosine domains, one which is important for receptor tyrosine kinase activity and eight which interact with different signal transduction proteins. The latter include the growth factor receptor binding protein Grb2, phosphatidylinositol 3-kinase, phospholipase C-gamma, the GTPase-activating protein GAP, the Src tyrosine kinase, and protein tyrosine phosphatase 1D (56). We have characterized signaling pathways stemming from three of these PDGF receptor binding proteins (see below).

5. ROLE OF THE ERK SIGNALING PATHWAY IN CELL CYCLE PROGRESSION

Grb2 is found in a stable complex with the nucleotide exchange factor Son of sevenless (Sos). Binding of this complex to the receptor tyrosine kinase induces Sos to bind to Ras, a 21 kD membrane-bound GTPase. Once activated, Ras may activate a number of downstream signaling pathways, including the extracellular signal regulated kinase (ERK) pathway (figure 1). ERKs are members of the mitogen-activated protein (MAP) kinase superfamily of cytosolic serine/threonine kinases that participate in the transduction of growth and differentiation-promoting signals to the cell nucleus. ERK activation has been shown to be required for DNA synthesis in a wide variety of cell systems, including bovine, rat and human airway smooth muscle (45, 57-60). The classical route to ERK activation involves Ras, Raf-1, a 74 kD cytoplasmic serine/threonine kinase, and MAP kinase/ERK kinase kinase (MEK)-1, a 45 kD dual function kinase. Several reports confirm that these intermediates are involved in ERK activation in airway smooth muscle.

Airway myocyte cell cycle regulation




Figure 1. Model illustrating three signaling pathways regulating transcription from the cyclin D₁ promoter in airway smooth muscle. The ERK and Rac1/PI 3-kinase pathways positively regulate cyclin D₁ expression (pathways 1 and 2, respectively); a third pathway, the p38 pathway, negatively regulates transcription from the cyclin D₁ promoter. Potential pathways of cross-talk are shown by the broad grey arrows. MEKK refers to a specific MAP kinase kinase kinase called MAP kinase/ERK kinase; MKKK refers to a generic MAP kinase kinase kinase.

Microinjection of the anti-pan Ras neutralizing antibody inhibited DNA synthesis in human airway smooth muscle cells (61), and overexpression of a dominant-negative form of H-Ras inhibited PDGF-mediated ERK activation in bovine tracheal myocytes (62). (Interestingly, Ras did not appear to be necessary for phorbol ester-induced ERK activation.) Overexpression of a kinase-dead mutant of Raf-1 inhibited endothelin-induced ERK activation in rat tracheal myocytes (44). Finally, chemical or dominant-negative inhibition of MEK-1 inhibits ERK activation and DNA synthesis in bovine, rat and human airway smooth muscle cells (45, 57-60).

One point at which bovine tracheal myocyte signaling may diverge from other cell types concerns the activation of MEK-1. Activation of MEK-1 occurred independently of Raf-1 in these cells (63) and instead involves a novel MEK-1 kinase (64).

Although activated ERK has been demonstrated to induce phosphorylation or associate with numerous nuclear transcription factors, the precise downstream targets of ERK in airway smooth muscle cells are not known. However, as in other cell types (65-67), it has been demonstrated in bovine tracheal myocytes that ERK activation is an upstream activator of transcription from the cyclin D₁ promoter (68). Thus, the ERK pathway appears to constitute an important regulator of entry into the cell cycle and G1 progression in airway smooth muscle.

In NIH 3T3 cells, constitutive activation of MEK-1, while sufficient to increase ERK activation and expression of cyclin D₁, is insufficient for maximal phosphorylation of the retinoblastoma protein, degradation of the cyclin dependent kinase inhibitor p27, and cyclin A expression (69). Furthermore Ras, but not ERK was shown to be required for growth factor-induced degradation of p27 in IIC9 fibroblasts (70). Finally, ectopic overexpression of cyclin D₁ is insufficient for S-phase traversal (34, 36).

Taken together, these data suggest that the ERK/cyclin D₁ pathway is insufficient for cell cycle progression, and that Ras coordinates cell cycle progression by regulating signaling through both ERK-dependent and ERK-independent signaling pathways.

6. ROLE OF PI 3-KINASE

PI3-kinase is a heterodimeric lipid kinase comprised of an 85 kD regulatory subunit and a 110 kD catalytic subunit which phosphorylates phosphatidylinositol at the D-3 hydroxyl of the inositol ring, forming the phosphatidylinositides phosphatidylinositol 3-phosphate, phosphatidylinositol 3,4-diphosphate and phosphatidylinositol 3,4,5-triphosphate. As noted above, ligation of the PDGF receptor may induce binding and activation of PI 3-kinase (figure 1). Ras has also been noted to interact with PI 3-kinase (71). Growth factors activate PI 3-kinase in human (43, 72) and bovine airway smooth muscle cells (73). Chemical inhibitors of PI 3-kinase, wortmannin and LY294002, inhibit airway smooth muscle cyclin D₁ expression (73) and DNA synthesis (73-75). Overexpression of the catalytic subdomain of PI 3-kinase in bovine tracheal myocytes induced transcription from the cyclin D₁ promoter but failed to activate ERK (73), suggesting that PI 3-kinase signaling occurs independently of ERK. Similarly, wortmannin and LY-294002 markedly inhibited EGF-induced PI 3-kinase activation in human airway smooth muscle cells but had no effect on ERK activation (75).

7. ROLE OF RAC1

Phosphoinositide products of PI 3-kinase may also influence the translocation of guanine nucleotide exchange factors, the upstream activators of GTPases (figure 1) (76). The Rho family GTPases (Rho A-C, Rac1 and 2, and Cdc42), through their regulation of the actin cytoskeleton and interactions with multiple target proteins, may influence cell cycle progression. In Swiss 3T3

Airway myocyte cell cycle regulation

fibroblasts, Rac1 is required for cell cycle progression (77, 78). In addition, Rac1 (79) and Cdc42, but not RhoA (C. Bauerfeld, K. Page, M. Hershenson, unpublished data) have been shown to function as upstream activators of cyclin D₁ expression in bovine tracheal myocytes.

Rac1 forms part of the NADPH oxidase complex that generates reactive oxygen species such as H₂O₂. Intracellular reactive oxygen intermediates are increased following growth factor treatment of both bovine (79) and rat tracheal myocytes (80). Accordingly, treatment with antioxidants attenuates both growth factor induced cyclin D₁ expression and DNA synthesis in these cells (79, 80). Overexpression of active Rac1 does not activate ERK in bovine tracheal myocytes, and Rac1-induced transcription from the cyclin D₁ promoter is insensitive to the chemical MEK inhibitor PD98059 (79), suggesting that Rac1-mediated cell cycle progression, like that following activation of PI 3-kinase, is independent of ERK activity.

8. INHIBITION OF AIRWAY SMOOTH MUSCLE CELL PROLIFERATION

Substances which induce a sustained increase in the intracellular concentration of cyclic AMP (cAMP) have been long known to inhibit airway smooth muscle growth (52, 81-87). However, the precise mechanism by which this occurs is unknown. In bovine tracheal myocytes, pretreatment with forskolin decreases cyclin D₁ protein abundance and promoter activity while inducing the phosphorylation and DNA binding of cAMP response element binding protein (CREB)-1. Taken together, these data suggest that cAMP suppresses cyclin D₁ gene expression via phosphorylation and transactivation of CREB. However, it remains unclear whether this is the primary mechanism of cAMP-induced growth inhibition, or whether the inhibition of upstream signaling pathways is involved. cAMP does not inhibit growth factor-induced activation of ERKs in bovine (63) or rat (58) airway smooth muscle, suggesting that the effect of cAMP on growth does not involve the ERK signaling pathway. Glucocorticoids also inhibit human airway smooth muscle cyclin D₁ expression and DNA synthesis in an ERK-independent manner (88). On the other hand, PI 3-kinase (74) has been shown to be cAMP-sensitive.

Growth factor treatment of airway smooth muscle cells also induces activation of the two stress-activated MAP kinases, p38 and Jun amino-terminal kinase (JNK) (47, 62, 89), consistent with the notion that these intermediates, like ERKs, play a role in the growth regulation (figure 1). However, based on the other types of signals which activate p38 and JNK (cellular stress and proinflammatory cytokines), it is possible that these pathways are involved in growth inhibition, rather than mitogenesis (see below). The p38 MAP kinase family now consists of four isoforms. p38alpha was originally identified in lipopolysaccharide-stimulated mouse macrophages and was found to have substantial homology to the yeast high osmolarity glycerol kinase (90-93). Since then, three additional isoforms, beta, gamma and delta have been cloned (94-97). p38alpha, beta and delta are

somewhat ubiquitously expressed, whereas p38gamma is primarily restricted to skeletal muscle (96). The p38 MAP kinases are phosphorylated and activated by MAP kinase kinase (MKK)-3, MKK-4, and MKK-6. MKK-6 appears to strongly activate all p38 isoforms, whereas MKK-3 preferentially activates p38alpha and beta (98-102). MKK-4 appears to phosphorylate and activate both JNK1 and p38alpha (92, 93). A number of distinct MAP kinase kinase kinases have been found to activate MKK-3 and MKK-6, including MAP kinase/ERK kinase kinase (MEKK)-1 (103, 104), mixed lineage kinase (MLK)-2 (104) and MLK-3 (105), MAP three kinase (MTK)-1 (106), apoptosis signal-regulating kinase (ASK) (107) and TAK-1, a potential mediator of TGF-beta signaling (108).

We have recently obtained data suggesting that selective activation of p38 inhibits airway smooth muscle cell cycle progression (109), as it does in CCL39 hamster lung fibroblasts (67). Chemical inhibition of p38 by SB202190 and SB203580 increased transcription from the cyclin D₁ promoter and cyclin D₁ protein abundance. Furthermore, transient transfection of bovine myocytes with dominant-negative forms of MKK3 or MKK6 increased transcription from the cyclin D₁ promoter. On the other hand, overexpression of constitutively active mutants of MKK3 or MKK6 decreased both basal and PDGF-mediated cyclin D₁ promoter activity. Since cyclin D₁ is a critical regulator of cell cycle progression (39), these data suggest that p38 is an important negative regulator of cell cycle progression in airway smooth muscle cells.

Paradoxically, p38 may be stimulated by Ras and Rac1 (62, 110-112), suggesting that GTPases may simultaneously activate positive and negative growth regulatory pathways, perhaps as a safeguard against excessive growth.

9. POTENTIAL ROLE OF PROTEIN KINASE C (PKC) ISOFORMS

PKC is a complex family including three types of isoenzymes. The conventional isoforms (alpha, beta₁, beta₂ and gamma) are activated by calcium, phorbol esters and phosphatidylserine, whereas the novel isoforms (delta, epsilon, eta, theta and mu) are calcium-insensitive and activated by phorbol esters and phosphatidylserine. The atypical isoforms (zeta, tau/lambda) are calcium and phorbol ester-insensitive and activated by phosphatidylserine. PKC alpha, beta₁, beta₂, delta, epsilon, and zeta, but not gamma or eta, are expressed in bovine tracheal myocytes (113), whereas PKC alpha, beta₁, beta₂, delta, epsilon, theta, eta, zeta, tau and mu have each been identified in human tracheal myocytes (114). Conventional and novel PKCs may be activated *in vivo* by diacylglycerol that is formed from phospholipids upon receptor-mediated activation of phospholipases. As noted above, ligand binding of the PDGF receptor leads to the phosphorylation of tyrosine residues which then serve as binding sites for signal transduction molecules containing Src-homology 2 (SH2) domains, including phospholipase C-gamma. Accordingly, growth factor stimulation has been noted to activate PKC delta in vascular smooth muscle (115), NIH 3T3 cells and monocytes (116).

Airway myocyte cell cycle regulation

Different PKC isoforms may have distinct roles in the regulation of cell proliferation. For example, PKCzeta activity increases in proliferating human airway smooth muscle (114). In NIH3T3 cells, PKCepsilon is a powerful growth stimulus, whereas PKCalpha and delta inhibit growth (117). PKCdelta also inhibits cell cycle progression in vascular smooth muscle cells (118), capillary endothelial cells (119) and rat colonic epithelial cells (120). Finally, in preliminary studies, we have found that overexpression of active PKCdelta attenuates growth factor-induced transcription from the cyclin D₁ promoter, perhaps via the aforementioned p38 inhibitory pathway (K. Page, M. Hershenson, unpublished data).

10. PERSPECTIVE

In recent years, the signaling pathways regulating airway smooth muscle growth have been elucidated. Although the substances mitogenic for airway smooth muscle may vary across species lines, the signal transduction mechanisms linking receptor ligation with DNA synthesis appear to be highly conserved. For example, the ERK and PI 3-kinase signaling pathways appear to constitute the major paths required for cell proliferation in both human (60, 72, 121) and bovine airway smooth muscle cells (57, 73, 74).

The signaling pathways regulating cell proliferation appear to be comparable in vascular and airway smooth muscle. Activation of ERK and PI 3-kinase has been shown to be required for vascular smooth muscle proliferation (122-124), and inhibition of proliferation by cAMP occurs via suppression of cyclin D₁ expression (125, 126), as it does in airway smooth muscle (87). Finally, reactive oxygen species may be critical for vascular smooth muscle cell mitogenic signaling (127, 128), as they are in airway myocytes (79). Perhaps these species and tissue similarities are to be expected, as many aspects of MAP kinase cascades, GTPase signaling pathways and cell cycle regulation are highly conserved in eukaryotic species, including mammals, Drosophila, nematodes and yeast (129-135).

Data are lacking regarding the roles of cyclins and their inhibitors in lung diseases. As might be expected, expression of cyclin D₁ antisense RNA retards lung cancer growth (136, 137). In rodents exposed to hyperoxia, whole lung p21Cip1 accumulates during hyperoxic exposure (138), whereas after exposure, alveolar epithelial cell cyclin D₁ expression increases, coincident with proliferative repair (139). Finally, as noted above, bronchoalveolar lavage fluid from asthmatic airways increases cultured human airway smooth muscle cell cyclin D₁ protein abundance relative to that from control subjects (17). Elucidation of the signal transduction and cell cycle mechanisms regulating airway smooth muscle growth may provide insight into similar mechanisms that occur in the airways of patients with bronchopulmonary dysplasia and chronic severe asthma, and lead to therapeutic interventions.

11. ACKNOWLEDGEMENTS

These studies were supported by National Institutes of Health Grants HL54685, HL56399, HL63314 and a grant from the Blowitz-Ridgeway Foundation.

12. REFERENCES

1. M.P. Sparrow, M. Weichselbaum & P.B. McCray: Development of the innervation and airway smooth muscle in human fetal lung. *Am J Respir Cell Mol Biol* 20, 550-560 (1999)
2. A.A. Hislop & S.G. Haworth: Airway size and structure in the normal fetal and infant lung and the effect of premature delivery and artificial ventilation. *Am Rev Respir Dis* 140, 1717-26 (1989)
3. S.L. Sward-Comunelli, S.M. Mabry, W.E. Truog & D.W. Thibeault: Airway muscle in preterm infants: changes during development. *J Pediatr* 130, 570-576 (1997)
4. D.E. Johnson, B.A. Burke & W.R. Anderson: Alterations in airway epithelial and smooth muscle area in infants with bronchopulmonary dysplasia. *Am Rev Respir Dis* 141, A159 (1991)
5. N. Carroll, J. Elliot, A. Morton & A. James: The structure of large and small airways in nonfatal and fatal asthma. *Am Rev Respir Dis* 147, 405-10 (1993)
6. M.S. Dunhill, G.R. Massarella & J.A. Anderson: A comparison of the quantitative anatomy of the bronchi in normal subjects, in status asthmaticus, in chronic bronchitis and in emphysema. *Thorax* 24, 176-181 (1971)
7. T. Takizawa & W.M. Thurlbeck: Muscle and mucous gland size in the major bronchi of patients with chronic bronchitis, asthma and asthmatic bronchitis. *Am Rev Respir Dis* 104, 331-336 (1971)
8. B.E. Heard & S. Hossain: Hyperplasia of bronchial smooth muscle in asthma. *J Pathol* 110, 319-332 (1973)
9. R.E. Sobonya: Quantitative structural alterations in long-standing allergic asthma. *Am Rev Respir Dis* 130, 289-92 (1984)
10. A.L. James, P.D. Pare & J.C. Hogg: The mechanics of airway narrowing in asthma. *Am Rev Respir Dis* 139, 242-6 (1989)
11. M. Ebina, H. Yaegashi, R. Chiba, T. Takahashi, M. Motomiya & M. Tanemura: Hyperreactive site in the airway tree of asthmatic patients revealed by thickening of bronchial muscles. A morphometric study. *Am Rev Respir Dis* 141, 1327-32 (1990)
12. M. Saetta, A. Di Stefano, C. Rosina, G. Thiene & L.M. Fabbri: Quantitative structural analysis of peripheral airways and arteries in sudden fatal asthma. *Am Rev Respir Dis* 143, 138-43 (1991)
13. M. Ebina, T. Takahashi, T. Chiba & M. Motomiya: Cellular hypertrophy and hyperplasia of airway smooth muscles underlying bronchial asthma. A 3-D morphometric study. *Am Rev Respir Dis* 148, 720-6 (1993)
14. M.D. Cohen, V. Ciocca & R.A. Panettieri: Transforming growth factor-beta1 modulates human airway smooth muscle cell proliferation induced by mitogens. *Am J Respir Cell Mol Biol* 16, 85-90 (1997)
15. A.M. Vignola, P. Chanez, G. Chiappara, A. Merendino, E. Pace, A. Rizzo, A.M. laRocca, V. Bellia, G. Bonsignore & J. Bousquet: Transforming growth factor-beta expression in mucosa biopsies in asthma and chronic bronchitis. *Am J Respir Crit Care Med* 156, 591-599 (1997)
16. M. Amishima, M. Munakata, Y. Nasuhara, A. Sato, T. Takahashi, Y. Homma & Y. Kawakami: Expression of

Airway myocyte cell cycle regulation

epidermal growth factor and epidermal growth factor receptor immunoreactivity in the asthmatic human airway. *Am Rev Respir Crit Care Med* 157, 1907-1912 (1998)

17. E.T. Naureckas, I.M. Ndukwu, A.J. Halayko, C. Maxwell, M.B. Hershenson & J. Solway: Bronchoalveolar lavage fluid from asthmatic subjects is mitogenic for human airway smooth muscle. *Am Rev Respir Crit Care Med* 160, 2062-2066 (1999)

18. M.B. Hershenson, M.D. Kelleher, E.T. Naureckas, M.K. Abe, V.J. Rubinstein, A. Zimmermann, A.M. Bendele, J.A. McNulty, R.A. Panettieri & J. Solway: Hyperoxia increases airway cell S-phase traversal in immature rats in vivo. *Am J Respir Cell Mol Biol* 11, 296-303 (1994)

19. Z.L. Wang, B.A. Walker, T.D. Weir, M.C. Yarema, C.R. Roberts, M. Okazawa, P.D. Pare & T.R. Bai: Effect of chronic antigen and beta 2 agonist exposure on airway remodeling in guinea pigs. *Am Rev Respir Crit Care Med* 152, 2097-104 (1995)

20. R.A. Panettieri, R.K. Murray, A.J. Eszterhas, G. Bilgen & J.G. Martin: Repeated allergen inhalations induce DNA synthesis in airway smooth muscle and epithelial cells in vivo. *Am J Physiol* 274, L417-L424 (1998)

21. J.T. Winston & W.J. Pledger: Growth factor regulation of cyclin D1 mRNA expression through protein synthesis-dependent and -independent mechanisms. *Mol Cell Biol* 15, 1133-44 (1995)

22. J. Kato, M. Matsuoka, K. Polyak, J. Massague & C. Sherr: Cyclic AMP-induced G1 phase arrest mediated by an inhibitor (p27Kip1) of cdk4 activation. *Cell* 79, 487-496 (1994)

23. X. Yue, H. Zhang & D. Beach: D-type cyclins associate with multiple protein kinases and the DNA replication and repair factor PCNA. *Cell* 71, 505-514 (1992)

24. J.W. Harper, G.R. Adami, N. Wei, K. Keyomarsi & S.J. Elledge: The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75, 805-16 (1993)

25. R. Bravo, R. Frank, P.A. Blundell & H. Macdonald-Bravo: Cyclin/PCNA is the auxiliary protein of DNA polymerase-delta. *Nature* 326, 515-517 (1987)

26. M. Pagano, S.W. Tam, A.M. Theodoras, P. Beer-Romero, G. Del Sal, V. Chau, P.R. Yew, G.F. Draetta & M. Rolfe: Role of the ubiquitin-proteasome pathway in regulating abundance of the cyclin-dependent kinase inhibitor p27. *Science* 269, 682-5 (1995)

27. J.Y. Kato, M. Matsuoka, D.K. Strom & C.J. Sherr: Regulation of cyclin D-dependent kinase 4 (cdk4) by cdk4-activating kinase. *Mol Cell Biol* 14, 2713-21 (1994)

28. H. Zhang, G.J. Hannon & D. Beach: p21-containing cyclin kinases exist in both active and inactive states. *Genes Devel* 8, 1750-8 (1994)

29. J. Kato, H. Matsushime, S.W. Hiebert, M.E. Ewen & C.J. Sherr: Direct binding of cyclin D to the retinoblastoma gene product (pRb) and pRb phosphorylation by the cyclin D-dependent kinase CDK4. *Genes Devel* 7, 331-42 (1993)

30. J.W. Ludlow, J. Shon, J.M. Pipas, D.M. Livingston & J.A. DeCaprio: The retinoblastoma susceptibility gene product undergoes cell cycle-dependent dephosphorylation and binding to and release from SV40 large T. *Cell* 60, 387-396 (1990)

31. J.R. Nevins: E2F: a link between the Rb tumor suppressor protein and viral oncoproteins. *Science* 258, 424-9 (1992)

32. J. Winston, F. Dong & W.J. Pledger: Differential modulation of G1 cyclins and the Cdk inhibitor p27kip1 by platelet-derived growth factor and plasma factors in density-arrested fibroblasts. *J Biol Chem* 271, 11253-11260 (1996)

33. J. Rosenblatt, Y. Gu & D.O. Morgan: Human cyclin-dependent kinase 2 is activated during the S and G2 phases of the cell cycle and associates with cyclin A. *Proc Natl Acad Sci* 89, 2824-8 (1992)

34. D.E. Quelle, R.A. Ashmun, S.A. Shurtleff, J.Y. Kato, D. Bar-Sagi, M.F. Roussel & C.J. Sherr: Overexpression of mouse D-type cyclins accelerates G1 phase in rodent fibroblasts. *Genes Devel* 7, 1559-71 (1993)

35. M. Ohtsubo & J.M. Roberts: Cyclin-dependent regulation of G1 in mammalian fibroblasts. *Science* 259, 1908-12 (1993)

36. D. Resnitzky, M. Gossen, H. Bujard & S.I. Reed: Acceleration of the G1/S phase transition by expression of cyclins D1 and E with an inducible system. *Mol Cell Biol* 14, 1669-79 (1994)

37. J. Lukas, J. Bartkova, M. Rohde, M. Strauss & J. Bartek: Cyclin D1 is dispensable for G1 control in retinoblastoma gene-deficient cells independently of cdk4 activity. *Mol Cell Biol* 15, 2600-11 (1995)

38. M. Ohtsubo, A.M. Theodoras, J. Schumacher, J.M. Roberts & M. Pagano: Human cyclin E, a nuclear protein essential for the G1-to-S phase transition. *Mol Cell Biol* 15, 2612-24 (1995)

39. W. Xiong, R.G. Pestell, G. Watanabe, J. Li, M.R. Rosner & M.B. Hershenson: Cyclin D1 is required for S-phase traversal in bovine tracheal smooth muscle myocytes. *Am J Physiol* 272, L1205-L1210 (1997)

40. S.J. Hirst, P.J. Barnes & C.H. Twort: Quantifying proliferation of cultured human and rabbit airway smooth muscle cells in response to serum and platelet-derived growth factor. *Am J Respir Cell Mol Biol* 7, 574-81 (1992)

41. M.D. Kelleher, M.K. Abe, T.S. Chao, M. Jain, J.M. Green, J. Solway, M.R. Rosner & M.B. Hershenson: Role of MAP kinase activation in bovine tracheal smooth muscle mitogenesis. *Am J Physiol* 268, L894-901 (1995)

42. S.J. Hirst, P.J. Barnes & C.H.C. Twort: PDGF isoform-induced proliferation and receptor expression in human cultured airway smooth muscle cells. *Am J Physiol* 270, L415-428 (1996)

43. V.P. Krymskaya, Hoffman, R., Eszterhas, A., Kane, S., Ciocca, V., Panettieri, R.A. Jr: EGF activates ErbB-2 and stimulates phosphatidylinositol 3-kinase in human airway smooth muscle cells. *Am J Physiol* 276, L246-L255 (1999)

44. P. Vichi, A. Whelchel, H. Knot, M. Nelson, W. Kolch & J. Posada: Endothelin-stimulated ERK activation in airway smooth-muscle cells requires calcium influx and Raf activation. *Am Rev Respir Cell Mol Biol* 20, 99-105 (1999)

45. T.R. Walker, S.M. Moore, M.F. Lawson, R.A.J. Panettieri & E.R. Chilvers: Platelet-derived growth factor-BB and thrombin activate phosphoinositide 3-kinase and protein kinase B: role in mediating airway smooth muscle proliferation. *Mol Pharmacol* 54, 1007-15 (1998)

Airway myocyte cell cycle regulation

46. A. Whelchel, J. Evans & J. Posada: Inhibition of ERK activation attenuates endothelin-stimulated airway smooth muscle cell proliferation. *Am J Respir Cell Mol Biol* 16, 589-96 (1997)
47. P.S. Shapiro, J.N. Evans, R.J. Davis & J.A. Posada: The seven-transmembrane-spanning receptors for endothelin and thrombin cause proliferation of airway smooth muscle cells and activation of the extracellular regulated kinase and c-Jun NH₂-terminal kinase groups of mitogen-activated protein kinases. *J Biol Chem* 271, 5750-4 (1996)
48. K. Malarkey, E.R. Chilvers, M.F. Lawson & R. Plevin: Stimulation by endothelin-1 of mitogen-activated protein kinases and DNA synthesis in bovine tracheal smooth muscle cells. *Br J Pharmacol* 116, 2267-73 (1995)
49. A.G. Stewart, G. Grigoriadis & T. Harris: Mitogenic actions of endothelin-1 and epidermal growth factor in cultured airway smooth muscle. *Clin Exp Pharmacol Physiol* 21, 277-85 (1994)
50. R.A. Panettieri, Jr., I.P. Hall, C.S. Maki & R.K. Murray: alpha-Thrombin increases cytosolic calcium and induces human airway smooth muscle cell proliferation. *Am J Respir Cell Mol Biol* 13, 205-16 (1995)
51. R.A. Panettieri, P.A. Yadavish, A.M. Kelly, N.A. Rubinstein & M.I. Kotlikoff: Histamine stimulates proliferation of airway smooth muscle and induces c-fos expression. *Am J Physiol* 259, L365-71 (1990)
52. K. Maruno, A. Absood & S.I. Said: VIP inhibits basal and histamine-stimulated proliferation of human airway smooth muscle cells. *Am J Physiol* 268, L1047-51 (1995)
53. A. Ullrich & J. Schlessinger: Signal transduction by receptors with tyrosine kinase activity. *Cell* 61, 203-0 (1990)
54. M.A. Lemmon & J. Schlessinger: Regulation of signal transduction and signal diversity by receptor oligomerization. *Trends in Biol Sci* 19, 459-463 (1994)
55. M.F. White, S.E. Shoelson, H. Keutmann & C.R. Kahn: A cascade of tyrosine autophosphorylation in the beta-subunit activates the phosphotransferase of the insulin receptor. *J Biol Chem* 263, (1988)
56. L. Claesson-Welsh: Platelet-derived growth factor receptor signaling. *J Biol Chem* 269, 32023-32026 (1994)
57. A.Y. Karpova, M.K. Abe, J. Li, P. Liu, J.M. Rhee, W.L. Kuo & M.B. Hershenson: MEK1 is required for PDGF-induced ERK activation and DNA synthesis in tracheal myocytes. *Am J Physiol* 272, L558-L565 (1997)
58. A. Whelchel, J. Evans & J. Posada: Inhibition of ERK activation attenuates endothelin-stimulated airway smooth muscle proliferation. *Am J Respir Cell Mol Biol* 16, 589-596 (1997)
59. D.B. Lew, B.K. Dempsey, Y. Zhao, M. Muthalif, S. Fatima & K.U. Malik: beta-hexosaminidase-induced activation of p44/42 mitogen-activated protein kinase is dependent on p21Ras and protein kinase C and mediates bovine airway smooth-muscle proliferation. *Am Rev Respir Cell Mol Biol* 21, 111-118 (1999)
60. M.J. Orsini, V.P. Krymskaya, A.J. Eszterhas, J.L. Benovic, R.A. Panettieri & R.B. Penn: MAPK superfamily activation in human airway smooth muscle: mitogenesis requires prolonged p42/p44 activation. *Am J Physiol* 277, L479-L488 (1999)
61. A.J. Ammit, S.A. Kane & R.A.J. Panettieri: Activation of K-p21ras and N-p21ras, but not H-p21ras, is necessary for mitogen-induced airway smooth muscle proliferation. *Am J Respir Cell Mol Biol* (1999)
62. K. Page, J. Li & M.B. Hershenson: Platelet-derived growth factor stimulation of mitogen-activated protein kinases and cyclin D1 promoter activity in cultured airway smooth muscle cells: Role of Ras. *Am J Respir Cell Mol Biol* 20, (1999)
63. M.B. Hershenson, T.S. Chao, M.K. Abe, I. Gomes, M.D. Kelleher, J. Solway & M.R. Rosner: Histamine antagonizes serotonin and growth factor-induced mitogen-activated protein kinase activation in bovine tracheal smooth muscle cells. *J Biol Chem* 270, 19908-13 (1995)
64. S. Kartha, E.T. Naureckas, J. Li & M.B. Hershenson: Partial characterization of a novel mitogen-activated protein kinase/extracellular signal-regulated kinase activator in airway smooth-muscle cells. *Am J Respir Cell Mol Biol* 20, 1041-8 (1999)
65. C. Albanese, J. Johnson, G. Watanabe, N. Eklund, D. Vu, A. Arnold & R.G. Pestell: Transforming p21ras mutants and c-Ets-2 activate the cyclin D1 promoter through distinguishable regions. *J Biol Chem* 270, 23589-97 (1995)
66. G. Watanabe, A. Howe, R. Lee, C. Albanese, I. Shu, A. Karnezis, L. Zon, J. Kyriakis, K. Rundell & R. Pestell: Induction of cyclin D1 by simian virus 40 small T tumor antigen. *Proc Natl Acad Sci USA* 93, 12861-12866 (1996)
67. J. Lavoie, G. L'Allemain, A. Brunet, R. Muller & J. Pouyssegur: Cyclin D1 expression is regulated positively by p42/p44 MAPK and negatively by p38/HOGMAPK pathway. *J Biol Chem* 271, 20608-20616 (1996)
68. M. Ramakrishnan, N.L. Musa, J. Li, P. Liu, R.G. Pestell & M.B. Hershenson: Catalytic activation of extracellular signal-regulated kinases induces cyclin D1 expression in airway smooth muscle. *Am Rev Respir Cell Mol Biol* 18, 736-740 (1998)
69. M. Cheng, V. Sexl, C.J. Sherr & M.F. Roussel: Assembly of cyclin D-dependent kinase and titration of p27Kip1 regulated by mitogen-activated protein kinase kinase (MEK1). *Proc Natl Acad Sci* 95, 1091-6 (1998)
70. J.D. Weber, W. Hu, S.C. Jefcoat, Jr., D.M. Raben & J.J. Baldassare: Ras-stimulated extracellular signal-related kinase 1 and RhoA activities coordinate platelet-derived growth factor-induced G1 progression through the independent regulation of cyclin D1 and p27. *J Biol Chem* 272, 32966-71 (1997)
71. P. Rodriguez-Viciana, P.H. Warne, R. Dhand, B. Vanhaesebroeck, I. Gout, M.J. Fry, M.D. Waterfield & J. Downward: Phosphatidylinositol-3-OH kinase as a direct target of Ras. *Nature* 370, 527-32 (1994)
72. V.P. Krymskaya, R. Hoffman, A. Eszterhas, V. Ciocca & P.A. Panettieri: TGF-beta1 modulates EGF-stimulated phosphatidylinositol 3-kinase activity in human airway smooth muscle cells. *Am J Physiol* 273, L1220-L1227 (1997)
73. K. Page, J. Li, Y. Wang, S. Kartha & M.B. Hershenson: Regulation of cyclin D1 expression and DNA synthesis by phosphatidylinositol 3-kinase in airway smooth muscle cells. Submitted (1999)
74. P.H. Scott, C.M. Belham, J. al-Hafidh, E.R. Chilvers, A.J. Peacock, G.W. Gould & R. Plevin: A regulatory role for cAMP in phosphatidylinositol 3-kinase/p70 ribosomal S6 kinase-mediated DNA synthesis in platelet-derived-

Airway myocyte cell cycle regulation

growth-factor-stimulated bovine airway smooth-muscle cells. *Biochem J* 318, 965-71 (1996)

75. V.P. Krymskaya, R.B. Penn, M.J. Orsini, P.H. Scott, R.J. Plevin, T.R. Walker, A.J. Eszterhas, Y. Amrani, E.R. Chilvers & R.A.J. Panettieri: Phosphatidylinositol 3-kinase mediates mitogen-induced human airway smooth muscle cell proliferation. *Am J Physiol* 277, L65-L78 (1999)

76. M.A. Lemmon, K.M. Ferguson, R. O'Brien, P.B. Sigler & J. Schlessinger: Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin homology domain. *Proc Natl Acad Sci USA* 92, 10472-6 (1995)

77. M.F. Olson, A. Ashworth & A. Hall: An essential role for Rho, Rac, and Cdc42 GTPases in cell cycle progression through G1. *Science* 269, 1270-2 (1995)

78. N. Lamarche, N. Tapon, L. Stowers, P.D. Burbelo, P. Aspenstrom, T. Bridges, J. Chant & A. Hall: Rac and Cdc42 induce actin polymerization and G1 cell cycle progression independently of p65PAK and the JNK/SAPK MAP kinase cascade. *Cell* 87, 519-29 (1996)

79. K. Page, J. Li, J.A. Hodge, P.T. Liu, T.L. Vanden Hoek, L.B. Becker, R.G. Pestell, M.R. Rosner & M.B. Hershenson: Characterization of a Rac1 signaling pathway to cyclin D1 expression in airway smooth muscle cells. *J Biol Chem* 274, 22065-22071 (1999)

80. S.S. Brar, T.P. Kennedy, A.R. Whorton, T.M. Murphy, P. Chitano & J.R. Hoidal: Requirement for reactive oxygen species in serum-induced and platelet-derived growth factor-induced growth of airway smooth muscle. *J Biol Chem* 274, 20017-20026 (1999)

81. D.B. Lew, C. Nebigil & K.U. Malik: Dual regulation by cAMP of beta-hexosaminidase-induced mitogenesis in bovine tracheal myocytes. *Am J Respir Cell Mol Biol* 7, 614-9 (1992)

82. C. Florio, J.G. Martin, A. Styhler & S. Heisler: Antiproliferative effect of prostaglandin E2 in cultured guinea pig tracheal smooth muscle cells. *Am J Physiol* 266, L131-7 (1994)

83. J.P. Noveral & M.M. Grunstein: Adrenergic receptor-mediated regulation of cultured rabbit airway smooth muscle cell proliferation. *Am J Physiol* 267, L291-9 (1994)

84. P.R. Johnson, C.L. Armour, D. Carey & J.L. Black: Heparin and PGE2 inhibit DNA synthesis in human airway smooth muscle cells in culture. *Am J Physiol* 269, L514-9 (1995)

85. P.R. Tomlinson, J.W. Wilson & A.G. Stewart: Salbutamol inhibits the proliferation of human airway smooth muscle cells grown in culture: relationship to elevated cAMP levels. *Biochem Pharmacol* 49, 1809-19 (1995)

86. A.G. Stewart, P.R. Tomlinson & J.W. Wilson: beta2-adrenoreceptor agonist-mediate inhibition of human airway smooth muscle proliferation: importance of the duration of beta2-adrenoreceptor stimulation. *Br J Pharmacol* 121, 361-368 (1997)

87. N.L. Musa, M. Ramakrishnan, J. Li, P. Liu, R.G. Pestell & M.B. Hershenson: Forskolin inhibits cyclin D1 expression in airway smooth muscle. *Am Rev Respir Cell Mol Biol* 20, 352-358 (1999)

88. D. Fernandes, E. Guida, V. Koutsoubos, T. Harris, P. Vadiveloo, J.W. Wilson & A.G. Stewart: Glucocorticoids inhibit proliferation, cyclin D1 expression, and retinoblastoma protein phosphorylation, but not activity of

the extracellular-regulated kinases in human cultured airway smooth muscle. *Am J Respir Cell Mol Biol* 21, 77-88 (1999)

89. N.J. Pyne & S. Pyne: Platelet-derived growth factor activates a mammalian Ste20 coupled mitogen-activated protein kinase in airway smooth muscle. *Cell Signal* 9, 311-317 (1997)

90. J. Han, J.D. Lee, P.S. Tobias & R.J. Ulevitch: Endotoxin induces rapid protein tyrosine phosphorylation in 70Z/3 cells expressing CD14. *J Biol Chem* 268, 25009-14 (1993)

91. A. Minden, A. Lin, M. McMahon, C. Lange-Carter, B. Derijard, R.J. Davis, G.L. Johnson & M. Karin: Differential activation of ERK and JNK mitogen-activated protein kinases by Raf-1 and MEKK. *Science* 266, 1719-23 (1994)

92. A. Lin, A. Minden, H. Martinetto, F.X. Claret, C. Lange-Carter, F. Mercurio, G.L. Johnson & M. Karin: Identification of a dual specificity kinase that activates the Jun kinases and p38-Mpk2. *Science* 268, 286-90 (1995)

93. B. Derijard, J. Raingeaud, T. Barrett, I.H. Wu, J. Han, R.J. Ulevitch & R.J. Davis: Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. *Science* 267, 682-5 (1995)

94. Y. Jiang, C. Chen, Z. Li, W. Guo, J.A. Gegner, S. Lin & J. Han: Characterization of the structure and function of a new mitogen-activated protein kinase (p38beta). *J Biol Chem* 271, 17920-6 (1996)

95. Y. Jiang, H. Gram, M. Zhao, L. New, J. Gu, L. Feng, F. Di Padova, R.J. Ulevitch & J. Han: Characterization of the structure and function of the fourth member of p38 group mitogen-activated protein kinases, p38delta. *J Biol Chem* 272, 30122-8 (1997)

96. X.S. Wang, K. Diener, C.L. Manthey, S. Wang, B. Rosenzweig, J. Bray, J. Delaney, C.N. Cole, P.Y. Chan-Hui, N. Mantlo, H.S. Lichenstein, M. Zukowski & Z. Yao: Molecular cloning and characterization of a novel p38 mitogen-activated protein kinase. *J Biol Chem* 272, 23668-74 (1997)

97. B. Stein, M.X. Yang, D.B. Young, R. Janknecht, T. Hunter, B.W. Murray & M.S. Barbosa: p38-2, a novel mitogen-activated protein kinase with distinct properties. *J Biol Chem* 272, 19509-17 (1997)

98. J. Raingeaud, A.J. Whitmarsh, T. Barrett, B. Derijard & R.J. Davis: MKK3- and MKK6-regulated gene expression is mediated by the p38 mitogen-activated protein kinase signal transduction pathway. *Mol Cell Biol* 16, 1247-55 (1996)

99. C. Tournier, A.J. Whitmarsh, J. Cavanagh, T. Barrett & R.J. Davis: Mitogen-activated protein kinase kinase 7 is an activator of the c-Jun NH2-terminal kinase. *Proc Natl Acad Sci* 94, 7337-42 (1997)

100. X. Lu, S. Nemoto & A. Lin: Identification of c-Jun NH2-terminal protein kinase (JNK)-activating kinase 2 as an activator of JNK but not p38. *J Biol Chem* 272, 24751-4 (1997)

101. P.M. Holland, M. Suzanne, J.S. Campbell, S. Noselli & J.A. Cooper: MKK7 is a stress-activated mitogen-activated protein kinase kinase functionally related to hemipterous. *J Biol Chem* 272, 24994-8 (1997)

102. H. Enslen, J. Raingeaud & R.J. Davis: Selective activation of p38 mitogen-activated protein (MAP) kinase

Airway myocyte cell cycle regulation

- isoforms by the MAP kinase kinases MKK3 and MKK6. *J Biol Chem* 273, 1741-8 (1998)
103. S. Xu, D.J. Robbins, L.B. Christerson, J.M. English, C.A. Vanderbilt & M.H. Cobb: Cloning of rat MEK kinase 1 cDNA reveals an endogenous membrane-associated 195-kDa protein with a large regulatory domain. *Proc Natl Acad Sci USA* 93, 5291-5 (1996)
104. A. Cuenda & D.S. Dorow: Differential activation of stress-activated protein kinase kinases SKK4/MKK7 and SKK1/MKK4 by the mixed-lineage kinase-2 and mitogen-activated protein kinase kinase (MKK) kinase-1. *Biochem J* 333, 11-15 (1998)
105. L.A. Tibbles, Y.L. Ing, F. Kiefer, J. Chan, N. Iscove, J.R. Woodgett & N.J. Lassam: MLK-3 activates the SAPK/JNK and p38/RK pathways via SEK1 and MKK3/6. *EMBO J* 15, 7026-35 (1996)
106. M. Takekawa, F. Posas & H. Saito: A human homolog of the yeast Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates stress-induced activation of the p38 and JNK pathways. *EMBO J* 16, 4973-82 (1997)
107. H. Ichijo, E. Nishida, K. Irie, P. ten Dijke, M. Saitoh, T. Moriguchi, M. Takagi, K. Matsumoto, K. Miyazono & Y. Gotoh: Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* 275, 90-4 (1997)
108. K. Yamaguchi, K. Shirakabe, H. Shibuya, K. Irie, I. Oishi, N. Ueno, T. Taniguchi, E. Nishida & K. Matsumoto: Identification of a member of the MAPKKK family as a potential mediator of TGF-beta signal transduction. *Science* 270, 2008-11 (1995)
109. K. Page, J. Li, P. Liu, S. Kartha & M.B. Hershenson: Inhibition of p38 MAP kinase increases cyclin D1 expression in cultured airway smooth muscle cells. *Am Rev Respir Crit Care Med* 159, A532 (abstract) (1999)
110. A. Minden, A. Lin, T. Smeal, B. Derijard, M. Cobb, R. Davis & M. Karin: c-Jun N-terminal phosphorylation correlates with activation of the JNK subgroup but not the ERK subgroup of mitogen-activated protein kinases. *Mol Cell Biol* 14, 6683-8 (1994)
111. S. Zhang, J. Han, M.A. Sells, J. Chernoff, U.G. Knaus, R.J. Ulevitch & G.M. Bokoch: Rho family GTPases regulate p38 mitogen-activated protein kinase through the downstream mediator PAK1. *J Biol Chem* 270, 23934-6 (1995)
112. S. Bagrodia, B. Derijard, R.J. Davis & R.A. Cerione: Cdc42 and PAK-mediated signaling leads to Jun kinase and p38 mitogen-activated protein kinase activation. *J Biol Chem* 270, 27995-8 (1995)
113. B.L.J. Webb, M.A. Lindsay, P.J. Barnes & M.A. Giembycz: Protein kinase C isoforms in airway smooth muscle. *Biochem J* 324, 167-175 (1997)
114. S. Carlin, K.X. Yang, R. Donnelly & J.L. Black: Protein kinase C isoforms in human airway smooth muscle cells: activation of PKC during proliferation. *Am J Physiol* 276, L506-L512 (1999)
115. L. Leng, B. Du, S. Consigli & T.A. McCaffrey: Translocation of protein kinase C-delta by PDGF in cultured vascular smooth muscle cells: inhibition by TGF-beta 1. *Artery* 22, 140-154 (1996)
116. W. Li & J.H. Pierce: Protein kinase C delta, an important signaling molecule in the platelet-derived growth factor-beta receptor pathway. *Curr Top Microbiol Immunol* 211, 55-65 (1996)
117. P. Acs, Q.J. Wang, K. Bogi, A.M. Marquez, P.S. Lorenzo, T. Biro, Z. Szallasi, J.F. Muchinski & P.M. Blumberg: Both the catalytic and regulatory domains of protein kinase C chimeras modulate the proliferative properties of NIH3T3 cells. *J Biol Chem* 272, 28793-28799 (1997)
118. S. Fukumoto, Y. Nishizawa, M. Hosoi, H. Koyama, K. Yamakawa, S. Ohno & H. Morii: Protein kinase C inhibits the proliferation of vascular smooth muscle by suppressing G1 cyclin expression. *J Biol Chem* 272, 13816-13822 (1997)
119. E.O. Harrington, J. Loffler, P.R. Nelson, K.C. Kent, M. Simons & J.A. Ware: Enhancement of migration by protein kinase C alpha and inhibition of proliferation and cell cycle progression by protein kinase C delta in capillary endothelial cells. *J Biol Chem* 272, 7390-7397 (1997)
120. G.P. Perletti, E. Marras, P. Concaro, F. Piccinini & A.H. Tashjian: PKCdelta acts as a growth and tumor suppressor in rat colonic epithelial cells. *Oncogene* 18, 1251-1256 (1999)
121. T.R. Walker, R.A. Panettieri & E.R. Chilvers: Role of phosphoinositide 3-kinase activation in airway smooth muscle proliferation. *Am J Respir Crit Care Med* 155, A903 (abstract). (1997)
122. P.A. Wilden, Y.M. Agazie, R. Kaufman & S.P. Halenda: ATP-stimulated smooth muscle cell proliferation requires independent ERK and PI3K signaling pathways. *Am J Physiol* 275, H1209-H1215 (1998)
123. D. Bacqueville, F. Casagrande, B. Perret, H. Chap, J.M. Darbon & M. Breton-Douillon: Phosphatidylinositol 3-kinase inhibitors block aortic smooth muscle cell proliferation in mid-late G1 phase: effect on cyclin-dependent kinase 2 and the inhibitory protein p27KIP1. *Biochem Biophys Res Commun* 27, 630-636 (1998)
124. X.P. Xi, K. Graf, S. Goetze, E. Fleck, W.A. Hsueh & R.E. Law: Central role of the MAPK pathway in ang II-mediated DNA synthesis and migration in rat vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* 19, 73-82 (1999)
125. S. Fukumoto, H. Koyama, M. Hosoi, K. Yamakawa, S. Tanaka, H. Morii & Y. Nishizawa: Distinct role of cAMP and cGMP in the cell cycle control of vascular smooth muscle cells: cGMP delays cell cycle transition through suppression of cyclin D1 and cyclin-dependent kinase 4 activation. *Circ Res* 85, 985-991 (1999)
126. N. Kronemann, W.A. Nockher, R. Busse & V.B. Schini-Kerth: Growth-inhibitory effect of cyclic GMP- and cyclic AMP-dependent vasodilators on rat vascular smooth muscle cells: effect on cell cycle and cyclin expression. *Br J Pharmacol* 126, 349-357 (1999)
127. M. Sundaresan, Z.-X. Yu, V.J. Ferrans, K. Irani & T. Finkel: Requirement for generation of H₂O₂ for platelet-derived growth factor signal transduction. *Science* 270, 296-299 (1995)
128. G. Lu, E.L. Greene, T. Nagai & B.M. Egan: Reactive oxygen species are critical in the oleic acid-mediated mitogenic signaling pathway in vascular smooth muscle cells. *Hypertension* 32, 1003-1010 (1998)
129. I. Herskowitz: MAP kinase pathways in yeast: for mating and more. *Cell* 80, 187-97 (1995)
130. R. Treisman: Regulation of transcription by MAP kinase cascades. *Curr Opin Cell Biol* 8, 205-15 (1996)

Airway myocyte cell cycle regulation

131. A.J. Waskiewicz & J.A. Cooper: Mitogen and stress response pathways: MAP kinase cascades and phosphatase regulation in mammals and yeast. *Curr Opin Cell Biol* 7, 798-805 (1995)
132. N. Watanabe, P. Madaule, T. Reid, T. Ishizaki, G. Watanabe, A. Kakizuka, Y. Saito, K. Nakao, B.M. Jockusch & S. Narumiya: p140^{mdia}, a mammalian homolog of *Drosophila* diaphanous, is a target protein for Rho small GTPase and is a ligand for profilin. *EMBO J* 16, 3044-56 (1997)
133. G. Watanabe, Y. Saito, P. Madaule, T. Ishizaki, K. Fujisawa, N. Morii, H. Mukai, Y. Ono, A. Kakizuka & S. Narumiya: Protein kinase N (PKN) and PKN-related protein raphilin as targets of small GTPase Rho. *Science* 271, 645-8 (1996)
134. B.A. Edgar & C.F. Lehner: Developmental control of cell cycle regulators: a fly's perspective. *Science* 274, 1646-1652 (1996)
135. S.J. Elledge: Cell cycle checkpoints: preventing an identity crisis. *Science* 274, 1664-1672 (1996)
136. D.S. Schrump, A. Chen & U. Consoli: Inhibition of lung cancer proliferation by antisense cyclin D. *Cancer Gene Ther* 3, 131-135 (1996)
137. B. Driscoll, L. Wu, S. Buckley, F.L. Hall, K.D. Anderson & D. Warburton: Cyclin D1 antisense RNA destabilizes pRb and retards lung cancer cell growth. *Am J Physiol* 273, L941-L949 (1997)
138. M.A. O'Reilly, R.J. Staversky, R.H. Watkins & W.M. Maniscalco: Accumulation of p21(Cip1/WAF1) during hyperoxic lung injury in mice. *Am J Respir Cell Mol Biol* 19, 777-785 (1998)
139. K.C. Bui, S. Buckley, F. Wu, B. Uhal, I. Joshi, J. Liu, M. Hussain, I. Makhoul & D. Warburton: Induction of A- and D-type cyclins and cdc2 kinase activity during recovery from short-term hyperoxic lung injury. *Am J Physiol* 268, L625-35 (1995)

Key words: Asthma, Cyclin D₁, Extracellular Signal Regulated Kinase, P38 Mitogen-Activated Protein Kinase, Phosphatidylinositol 3-Kinase, Platelet-Derived Growth Factor, Rac, Ras, Signal Transduction, Review

Send correspondence to: Marc B. Hershenson, M.D., University of Chicago Children's Hospital, 5841 S. Maryland Ave., MC4064, Chicago, IL 60637-1470, Tel:773-702-9659, Fax: 773-702-4041, E-mail: mhershen@midway.uchicago.edu

This manuscript is available on line at:

<http://www.bioscience.org/2000/v5/d/page/fulltext.htm>